

Japan Charged-Particle Nuclear Reaction Data Group

Division of Physics, Graduate School of Science

Hokkaido University

060-0810 Sapporo, JAPAN

Telephone +81(JPN)-11-706-2684

Facsimile +81(JPN)-11-706-4850

E-mail nrdf@jcprg.sci.hokudai.ac.jp

Memo CP-E/029

Date: December 10, 2003
To: Distribution
From: OTSUKA Naohiko and KATŌ Kiyoshi
Subject: Double differential cross section integrated over partial angular range

We are compiling a spectroscopic study of hypernucleus by the (π^+ , K^+) reaction (T. Hasegawa *et al.*, Phys. Rev. Lett. **74**(1995)224). In this work, authors give double differential cross sections with respect to emission angle of positive kaon and excitation energy of residual nucleus.

Similar quantity is also found in T. Teranishi *et al.*, Phys. Lett. **B4074**(1997)110, where double differential cross sections with respect to emission angle and excitation energy of ^{11}Be are measured for $^1\text{H}(^{11}\text{Li}, n)^{11}\text{Be}$ reaction. We propose the following codes for both cases:

Dictionary 36 (Quantities)

, DA/DE, , IPA	Double-differential cross section integrated partial angular range)
, DA/DE, RSD, IPA	Double-differential cross section $d^2/dA(\text{Resid.})dE(\text{Resid.})$

Distribution:

J.H. Chang, KAERI	M. Chiba, JCPRG	F.E. Chukreev, CAJaD	S. Dunaeva, NDS
O. Gritzay, KINR	J. Hasegawa, JAERI	A. Kaltchenko, KINR	K. Kato, JCPRG
M. Kellett, NEADB	M. Lammer, NDS	S. Maev, CJD	V.N. Manokhin, CJD
V. McLane, NNDC	C. Nordborg, NEADB	P. Oblozinsky, NNDC	Y. Ohbayasi, JCPRG
N. Otsuka, JCPRG	V. Pronyaev, NDS	O. Schwerer, NDS	S. Takacs, ATOMKI.
S. Taova, VNIIEF	T. Tárkányi, ATOMKI	V. Varlamov, CDFE	M. Vlasov, KINR
M. Wirtz, NDS	V. Zerkín, NDS	Zhuang Y.X., CNDC	

Sample of coded entry (E1601.002):

SUBENT	E1601001	20031211	E160100100001
BIB	10	38	E160100100002
...			
EN-SEC	ANG-MIN is minimum polar angle between beam and positive kaon in laboratory system		E160100100020
	ANG-MAX is maximum polar angle between beam and positive kaon in laboratory system		E160100100021
			E160100100022
			E160100100023
ENDBIB	38	0	E160100100041
COMMON	3	3	E160100100042
ANG-MIN	ANG-MAX	MOM	E160100100043
ADEG	ADEG	GEV/C	E160100100044
2.0	14.0	1.06	E160100100045
ENDCOMMON	3	0	E160100100046
ENDSUBENT	45	0	E160100199999
SUBENT	E1601002	20031211	E160100200001
BIB	5	5	E160100200002
REACTION	(6-C-12(PIP,X)1-KP-0,,DA/DE,,IPA)		E160100200003
PART-DET	(KP)		E160100200004
EN-SEC	E-EXC: Excitation energy of 12(Lambda)C		E160100200005
ANALYSIS	(INTAD)integrated from 2 to 14 deg		E160100200006
STATUS	(CURVE)Data scanned from Fig.2, p225 in reference		E160100200007
ENDBIB	5	0	E160100200008
NOCOMMON	0	0	E160100200009
DATA	3	99	E160100200010
E-EXC	DATA	DATA-ERR	E160100200011
MEV	MU-B/MEV	MU-B/MEV	E160100200012
-1.366E+01	1.110E-05		E160100200013
-1.317E+01	8.896E-04		E160100200014
...			
	3.527E+01	2.840E-01 2.322E-02	E160100200110
	3.570E+01	2.350E-01 2.064E-02	E160100200111
ENDDATA	101	0	E160100200112
ENDSUBENT	111	0	E160100299999

Sample of coded entry (E1670.002):

SUBENT	E1670002	20031211			E167000200001
BIB	8	31			E167000200002
REACTION	(1-H-1(3-LI-11,N)4-BE-11,,DA/DE,RSD,IPA/RAW)				E167000200003
	DATA: Uncorrected for detector acceptance				E167000200004
PART-DET	(P)				E167000200005
	(N)				E167000200006
	(3-LI-9)				E167000200007
EN-SEC	ANG-MIN is polar angle between beam and 11Be in laboratory system				E167000200008
	ANG-MAX is polar angle between beam and 11Be in laboratory system				E167000200009
	(E-EXC,4-BE-11)Excitation energy of 11Be. This energy is calculated by $E(d) + 20.14$ MeV where $E(d)$ is decay energy defined in p112 of the reference and 20.14 MeV is threshold energy for the $9\text{Li} + p + n$ channel [G.Audi, A.H.Wapstra, Nucl. Phys. A 595(1995)409].				E167000200012
MISC-COL	(MISC1)Decay energy $E(d)$ defined in p112 of the reference				E167000200019
	(MISC2)Detector acceptance. True energy spectrum can be obtained by dividing tabulated data by this acceptance.				E167000200021
SAMPLE	Chemical-form of target is CH ₂ . (CH ₂) _n and C target are used, C target was used to subtract the contributions of carbon nuclei in the (CH ₂) _n target. Target-thickness: 191 and 188 mg/cm ² for (CH ₂) _n and C target, respectively.				E167000200024
METHOD	(COINC)Detect 9Li, proton and neutron in coincidence.				E167000200029
ERR-ANALYS	(ERR-1)Mainly due to the uncertainty in neutron detection efficiency				E167000200030
STATUS	(TABLE)Data (Fig.2(a), p112 in reference) received by e-mail from T.Teranishi (2003.09.17)				E167000200032
ENDBIB	31	0			E167000200034
COMMON	3	3			E167000200035
ANG-MIN	ANG-MAX	ERR-1			E167000200036
ADEG	ADEG	PER-CENT			E167000200037
0.0	5.0	10.0			E167000200038
ENDCOMMON	3	0			E167000200039
DATA	5	100			E167000200040
E-EXC	MISC1	DATA	ERR-S	MISC2	E167000200041
MEV	MEV	MB/MEV	MB/MEV	NO-DIM	E167000200042
20.19	0.05	0.0	0.0	0.03	E167000200043
20.29	0.15	0.02	0.01	0.095	E167000200044
...					
29.99	9.85	0.03	0.02	0.071	E167000200141
30.09	9.95	0.02	0.02	0.069	E167000200142
ENDDATA	102	0			E167000200143
ENDSUBENT	142	0			E167000299999