

**NATIONAL NUCLEAR DATA CENTER
Bldg. 197D
Brookhaven National Laboratory
P. O. Box 5000
Upton, NY 11973-5000 U.S.A.**

(Internet) "NNDC@BNL.GOV"

Telephone: (516)344-2902
FAX: (516)344-2806

Memo CP-C/254

DATE: August 26, 1999
TO: Distribution
FROM: V. McLane
SUBJECT: Resonance strength

Please make the following dictionary updates.

Add to Dictionary (Reports)

STR Strength

Add to Dictionary 36 (Quantities)

,WID/STR Resonance strength

A proposed LEXFOR entry is attached

Distribution

M. Chiba, Sapporo

F. E. Chukreev, CaJaD

S. Dunaeva, Sarov

K. Kato, JCPDG

M. Kellett, NEADB

V. N. Manokhin, CJD

S. Maev, CJD

O. Schwerer, NDS

S. Tak<cs, ATOMKI

F. T. Tárkányi, ATOMKI

Y. Tendow, RIKEN

V. Varlamov, CDFE

Zhuang Youxiang, CNDC

NNDC (3)

LEXFOR (proposed)

Resonance Strength

The resonance strength is defined as:

$$\omega\Gamma = \frac{2J+1}{(2j_i+1)(2j_p+1)} \frac{\Gamma_i\Gamma_r}{\Gamma}$$

Where:

J = spin of resonance,

j_p = spin of incident projectile,

j_t = spin of target,

ε_p = partial width for formation of resonance by incident particle p,

ε_r = partial width for decay of resonance by reaction channel r,

ε = total width of resonance.

Resonance strengths are determined experimentally by measuring the area under the resonant yield curve:

$$\omega\Gamma = \frac{2\varepsilon}{\lambda_R} \frac{A_t}{A_t + A_p} Y_r$$

where: λ_R = particle wavelength at the resonance energy,

γ = stopping power

EXFOR coding

REACTION string: ,WID/STR

Units: energy, e.g., EV

The energy coded is the resonance energy as for other resonance data.